

STEVAN DOUGLAS LOONEY

Attorney at Law

[REDACTED]
Albuquerque, NM 87114

January 3, 2023

CERTIFIED MAIL-RETURN RECEIPT REQUESTED

Raul Torrez
Attorney General State of New Mexico
408 Galisteo Street
Villagra Building
Santa Fe, NM 87501

Re: Urgent Request to the Office of the Attorney General of the State of New Mexico to Enforce the Laws of the State of New Mexico to Protect the Health, Safety and Welfare of the Citizens of New Mexico

Dear Attorney General Torrez:

Numerous statutory civil laws have been and are being violated and crimes of unimaginable dimensions are being committed every day within the borders of the State of New Mexico. These civil violations and criminal activities began prior to 2020 and increased in frequency, intensity and proportion in 2021. They continue to this very day, not only unabated, but also unchallenged, uninvestigated and unpunished. As a result, the dire consequences to the citizens of New Mexico continue and cannot be overstated in their impact, nor underestimated in their breath, scope and severity. Sadly, these consequences include not only yet uncalculated serious physical and psychological injuries, but also a multitude of deaths.

Certain violations of New Mexico civil, if not criminal, laws were brought to the attention of your predecessor, Hector Balderas, during the time the citizens of this State reposed their trust and confidence in him when he occupied the Office of Attorney General. I know because, in July of 2021, I was one such person who reached out and implored him to discharge his duties and exercise the power and authority he held as the Attorney General to, at a minimum, launch civil investigative demands into the conduct of the perpetrators of the civil wrongs and very likely criminal conduct.

I have with this letter provided you with a copy of the July 23, 2021, letter (and enclosures) that I authored and personally mailed to Mr. Balderas, certified-returned receipt requested. While the letter was received by Mr. Balderas and the Office of the Attorney General, no response at all to

the letter was received by me. I pray that this letter does not likewise fall on uncaring or deaf ears, motivated more by political and partisan considerations, than by noble considerations such as integrity, compassion, humanity and, equally so, the rule of law.

By now you may have determined that the civil wrongs and criminal acts I am referring to concern SARS-CoV-2, masquerading under the moniker “Covid 19 disease”, and the experimental, unapproved, emergency use authorized (“EUA”), mRNA gene therapies which are deceptively and unconscionably called vaccines. These “vaccines” are the product of Pfizer/Bio-N-Tech and Moderna, among others, and were produced and marketed by these companies in the ordinary course of their business operations and made available to the citizens of New Mexico who, in turn, had these products injected into their bodies by deceptive and fraudulent means and methods, and without their informed consent.

I respectfully urge you to read and understand the letter and the attachments I presented to Mr. Balderas in July of 2021 and which I now also present to you. However, much has happened since July of 2021 regarding these so-called vaccines and their myriad adverse events. And none of it is good. For instance, by July of 2021 many tens of thousands of documented adverse events from these “vaccines” had been reported to the Vaccine Adverse Events Reporting System (“VAERS”). More disturbingly, according to the Centers for Disease Control and Prevention as of July 19, 2021, 12,313 deaths caused by these products had been reported to VAERS. As of December 2022, the adverse events, including deaths, is many, many times the number reported to VAERS as of July of 2021. According to a Harvard study of the VAERS, for various reasons, including that VAERS is a voluntary reporting system, adverse events, including deaths, are under reported by a factor of nearly 99%. That is, for every adverse event reported to VAERS, nearly 99 adverse events go unreported.

What has transpired with these experimental injections is demonstrably and provably inimical to the health, safety and welfare of the citizens of New Mexico, to say nothing of people elsewhere. The personal and emotional injury, as well as the death toll, from these so-called vaccines has increased dramatically and continues to increase every day. For example, while Pfizer/Bio-N-Tech initially disclosed to the public only relatively mild adverse side effects from their product, such as injection site pain, fatigue, headache, nausea, muscle pain, chills, and ‘feeling unwell’, these companies knew, but did not voluntarily publicly disclose, that through February 28, 2021, nearly 1,000 separate categories or types of “Adverse Events of Special Interest” (“AESI”) had been identified and reported. To establish that fact, I have attached a document as **Exhibit 1** prepared by or for Pfizer titled “*5.3.6 Cumulative Analysis of Post-Authorization Adverse Event Reports of PF-07302048 (BNT162B2) Received Through 28-Feb-2021*”. Attached to that document is another document titled “*Appendix 1. List of Adverse Events of Special Interest*”. That list contains approximately 1,000 AESIs that were known to Pfizer (and the FDA), but were neither voluntarily nor timely disclosed to the public. In fact, Pfizer and the FDA sought to withhold and hide this information and these AESIs from public view for 70-75 years. It was not until early 2022 when a federal court ordered the FDA to produce some 50,000 pages of documents related to the EUA of Pfizer’s BNT162B2, that is, Pfizer’s experimental mRNA “vaccine”, that these

AESIs were publicly disclosed. Indeed, the documents Pfizer was forced to disclose by a federal court order also showed that by February 28, 2021, not long after the release of these “vaccines” into the market, so many AESIs were being reported that Pfizer had to hire 1,800 to 2,400 additional full-time employees, on top of the existing 800 full-time Pfizer employees, just to process the AESIs.

Increasing numbers of people, including millions of Americans, recognize and experience the dangers of these products at a personal level. One measure of this recognition is the under-reported number of adverse events reported to VAERS. Peer reviewed studies conducted using the available public data also attest to the serious AESIs following mRNA COVID-19 vaccinations. See, e.g., <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9428332/> Another measure of the growing awareness of the ASEIs is found in a very recent poll conducted by Rasmussen Reports. Among other things, the results of that poll show that 28% of American adults, Democrats and Republicans alike, know, or believe they know, someone who died as a result of being injected with these products. These are deaths not merely AESIs that did not result in death. See, https://www.rasmussenreports.com/public_content/politics/public_surveys/died_suddenly_more_than_1_in_4_think_someone_they_know_died_from_covid_19_vaccines Yet, unlike other products that caused just a few deaths and were withdrawn from the market (including the Swine Flu vaccine and the Rotavirus vaccine Rotashield), these mRNA experimental gene therapies are not withdrawn from the market, rather they are aggressively promoted at every level and pushed on everyone incessantly.

What I have presented above is just a small sample of the carnage, fraud and deception undertaken and still underway, and touches upon only a fraction of the injuries including, myocarditis and pericarditis (serious heart conditions), and very direly, irreparable damage to the reproductive systems of women and men alike, and the deaths that have directly and proximately occurred due to the introduction of these products into the stream of commerce and the blood stream and organs of many millions of human beings. The deception, fraud and resulting injury perpetrated by the purveyors of these toxic and deleterious products will continue unless and until people, people such as you, with the power and authority to stop the carnage stand up and do so. You will not stand alone in this effort. The Attorneys General of states such as Louisiana and Missouri have already boldly stepped forward to protect the citizens of their states. So, too, the state of Florida. The Supreme Court of Florida has recently approved empaneling a Grand Jury to investigate the introduction of these products into Florida, including investigating their safety, efficacy and necessity, or the lack thereof. In addition, many hundreds, if not thousands, of medical doctors, virologist, immunologist and other men and women of medicine and science in the USA and around the world are in ever-increasing numbers coming forward to expose both the absence of proper medical practices and good science surrounding these products and the dangers they pose to the health, safety and welfare of human beings who are injected with these toxic products.

With the foregoing in mind, I respectfully direct your attention to several civil statutes promulgated by the New Mexico Legislature that not only empower you to, but often mandate that you take all deliberate action—including initiating civil investigative demands—to protect the health, safety

and welfare of all New Mexicans from products such as these experimental, mRNA gene therapies falsely being called “vaccines”.

The New Mexico civil statutes that I have identified to date are as follows:

- The New Mexico Unfair Practices Act, NMSA, Sections 57-12-1 *et seq.*
- The Fraud Against Taxpayers Act, NMSA, Sections 44-9-1 *et seq.*
- The Anti-Trust Act (Restraint of Trade), NMSA, Sections 57-1-1 *et seq.*
- The False Advertising Act, NMSA, Sections 57-15-1 *et seq.*
- The Vaccine Purchasing Act, NMSA, Sections 24-5A-1 *et seq.*
- The Drug and Cosmetics Act, NMSA, Sections 26-1-1 *et seq.*
- The Public Health Emergency Response Act, NMSA, Sections 12-10A-1 *et seq.*
- The Public Health Act, NMSA, Sections 24-1-1 *et seq.*

Additionally, federal statutes such as the RICO Act have been violated, as well as New Mexico criminal laws.

As the Attorney General of New Mexico you not only have the power and authority, but the duty, to launch investigations to determine not just whether but which civil and criminal statutes have been violated by the producers and purveyors of these products. Civil investigative demands are but one important preliminary tool available to you in pursuit of this necessary and noble cause.

On behalf of the citizens of New Mexico I implore you to promptly file the appropriate pleadings and take all necessary steps in the courts of competent jurisdiction to first obtain temporary restraining orders and then permanent injunctions to enjoin the unlawful conduct of these companies, and to seek and obtain damages, including treble and punitive damages, as well as restitution for those many thousands of New Mexicans injured or killed by these products. And, as to the crimes that have been committed, to identify those crimes and then file criminal complaints or petition the appropriate court to empanel a Grand Jury to seek indictments against the researchers, developers, manufacturers and distributors of these so-called vaccines.

The courtesy of a substantive response to the issues raised in this letter is requested. I am available and happy to discuss the substance of this letter with you should you wish to do so.

Respectfully,



Stevan Douglas Looney

SDL/sdl
Enclosure

5.3.6 CUMULATIVE ANALYSIS OF POST-AUTHORIZATION ADVERSE EVENT REPORTS OF PF-07302048 (BNT162B2) RECEIVED THROUGH 28-FEB-2021

Report Prepared by:

Worldwide Safety

Pfizer

The information contained in this document is proprietary and confidential. Any disclosure, reproduction, distribution, or other dissemination of this information outside of Pfizer, its Affiliates, its Licensees, or Regulatory Agencies is strictly prohibited. Except as may be otherwise agreed to in writing, by accepting or reviewing these materials, you agree to hold such information in confidence and not to disclose it to others (except where required by applicable law), nor to use it for unauthorized purposes.

ADVERSE EVENTS KNOWN TO PFIZER AND THE FDA BEFORE, ON AND AFTER FEBRUARY 28, 2021 THAT PFIZER AND THE FDA DID NOT WANT DISCLOSED TO THE PUBLIC BUT WHICH THEY WANTED BURIED FROM PUBLIC VIEW AND DISCLOSURE FOR 70-75 YEARS. THESE ADVERSE EVENTS WERE MADE KNOWN TO THE PUBLIC IN MARCH OF 2022 ONLY BECAUSE A FEDERAL COURT JUDGE ORDERED THAT THEY BE RELEASED TO THE PUBLIC. PER THE ORDER THE FDA IS REQUIRED TO PRODUCE ADDITIONAL DOCUMENTS.

090177e196ea1800ApprovedApproved On: 30-Apr-2021 09:26 (GMT)



Appendix 1
Attached

3.1.3. Review of Adverse Events of Special Interest (AESIs)

Please refer to Appendix 1 for the list of the company’s AESIs for BNT162b2.

The company’s AESI list takes into consideration the lists of AESIs from the following expert groups and regulatory authorities: Brighton Collaboration (SPEAC), ACCESS protocol, US CDC (preliminary list of AESI for VAERS surveillance), MHRA (unpublished guideline).

The AESI terms are incorporated into a TME list and include events of interest due to their association with severe COVID-19 and events of interest for vaccines in general.

The AESI list is comprised of MedDRA PTs, HLTs, HLTs or MedDRA SMQs and can be changed as appropriate based on the evolving safety profile of the vaccine.

Table 7 provides a summary review of cumulative cases within AESI categories in the Pfizer safety database. This is distinct from safety signal evaluations which are conducted and included, as appropriate, in the Summary Monthly Safety Reports submitted regularly to the FDA and other Health Authorities.

Table 7. AESIs Evaluation for BNT162b2

AESIs ^a Category	Post-Marketing Cases Evaluation ^b Total Number of Cases (N=42086)
<p>Anaphylactic Reactions <i>Search criteria: Anaphylactic reaction SMQ (Narrow and Broad, with the algorithm applied), selecting relevant cases according to BC criteria</i></p>	<p>Please refer to the Risk ‘Anaphylaxis’ included above in Table 4.</p>
<p>Cardiovascular AESIs <i>Search criteria: PTs Acute myocardial infarction; Arrhythmia; Cardiac failure; Cardiac failure acute; Cardiogenic shock; Coronary artery disease; Myocardial infarction; Postural orthostatic tachycardia syndrome; Stress cardiomyopathy; Tachycardia</i></p>	<ul style="list-style-type: none"> • Number of cases: 1403 (3.3% of the total PM dataset), of which 241 are medically confirmed and 1162 are non-medically confirmed; • Country of incidence: UK (268), US (233), Mexico (196), Italy (141), France (128), Germany (102), Spain (46), Greece (45), Portugal (37), Sweden (20), Ireland (17), Poland (16), Israel (13), Austria, Romania and Finland (12 each), Netherlands (11), Belgium and Norway (10 each), Czech Republic (9), Hungary and Canada (8 each), Croatia and Denmark (7 each), Iceland (5); the remaining 30 cases were distributed among 13 other countries; • Subjects’ gender: female (1076), male (291) and unknown (36); • Subjects’ age group (n = 1346): Adult^c (1078), Elderly^d (266) Child^e and Adolescent^f (1 each); • Number of relevant events: 1441, of which 946 serious, 495 non-serious; in the cases reporting relevant serious events; • Reported relevant PTs: Tachycardia (1098), Arrhythmia (102), Myocardial infarction (89), Cardiac failure (80), Acute myocardial infarction (41), Cardiac failure acute (11), Cardiogenic shock and Postural orthostatic tachycardia syndrome (7 each) and Coronary artery disease (6); • Relevant event onset latency (n = 1209): Range from <24 hours to 21 days, median <24 hours;

090177e196ea1800\Approved\Approved On: 30-Apr-2021 09:26 (GMT)

APPENDIX 1. LIST OF ADVERSE EVENTS OF SPECIAL INTEREST

1p36 deletion syndrome; 2-Hydroxyglutaric aciduria; 5'nucleotidase increased; Acoustic neuritis; Acquired C1 inhibitor deficiency; Acquired epidermolysis bullosa; Acquired epileptic aphasia; Acute cutaneous lupus erythematosus; Acute disseminated encephalomyelitis; Acute encephalitis with refractory, repetitive partial seizures; Acute febrile neutrophilic dermatosis; Acute flaccid myelitis; Acute haemorrhagic leukoencephalitis; Acute haemorrhagic oedema of infancy; Acute kidney injury; Acute macular outer retinopathy; Acute motor axonal neuropathy; Acute motor-sensory axonal neuropathy; Acute myocardial infarction; Acute respiratory distress syndrome; Acute respiratory failure; Addison's disease; Administration site thrombosis; Administration site vasculitis; Adrenal thrombosis; Adverse event following immunisation; Ageusia; Agranulocytosis; Air embolism; Alanine aminotransferase abnormal; Alanine aminotransferase increased; Alcoholic seizure; Allergic bronchopulmonary mycosis; Allergic oedema; Alloimmune hepatitis; Alopecia areata; Alpers disease; Alveolar proteinosis; Ammonia abnormal; Ammonia increased; Amniotic cavity infection; Amygdalohippocampectomy; Amyloid arthropathy; Amyloidosis; Amyloidosis senile; Anaphylactic reaction; Anaphylactic shock; Anaphylactic transfusion reaction; Anaphylactoid reaction; Anaphylactoid shock; Anaphylactoid syndrome of pregnancy; Angioedema; Angiopathic neuropathy; Ankylosing spondylitis; Anosmia; Antiacetylcholine receptor antibody positive; Anti-actin antibody positive; Anti-aquaporin-4 antibody positive; Anti-basal ganglia antibody positive; Anti-cyclic citrullinated peptide antibody positive; Anti-epithelial antibody positive; Anti-erythrocyte antibody positive; Anti-exosome complex antibody positive; Anti-GAD antibody negative; Anti-GAD antibody positive; Anti-ganglioside antibody positive; Antigliadin antibody positive; Anti-glomerular basement membrane antibody positive; Anti-glomerular basement membrane disease; Anti-glycyl-tRNA synthetase antibody positive; Anti-HLA antibody test positive; Anti-IA2 antibody positive; Anti-insulin antibody increased; Anti-insulin antibody positive; Anti-insulin receptor antibody increased; Anti-insulin receptor antibody positive; Anti-interferon antibody negative; Anti-interferon antibody positive; Anti-islet cell antibody positive; Antimitochondrial antibody positive; Anti-muscle specific kinase antibody positive; Anti-myelin-associated glycoprotein antibodies positive; Anti-myelin-associated glycoprotein associated polyneuropathy; Antimyocardial antibody positive; Anti-neuronal antibody positive; Antineutrophil cytoplasmic antibody increased; Antineutrophil cytoplasmic antibody positive; Anti-neutrophil cytoplasmic antibody positive vasculitis; Anti-NMDA antibody positive; Antinuclear antibody increased; Antinuclear antibody positive; Antiphospholipid antibodies positive; Antiphospholipid syndrome; Anti-platelet antibody positive; Anti-prothrombin antibody positive; Antiribosomal P antibody positive; Anti-RNA polymerase III antibody positive; Anti-saccharomyces cerevisiae antibody test positive; Anti-sperm antibody positive; Anti-SRP antibody positive; Antisynthetase syndrome; Anti-thyroid antibody positive; Anti-transglutaminase antibody increased; Anti-VGCC antibody positive; Anti-VGKC antibody positive; Anti-vimentin antibody positive; Antiviral prophylaxis; Antiviral treatment; Anti-zinc transporter 8 antibody positive; Aortic embolus; Aortic thrombosis; Aortitis; Aplasia pure red cell; Aplastic anaemia; Application site thrombosis; Application site vasculitis; Arrhythmia; Arterial bypass occlusion; Arterial bypass thrombosis; Arterial thrombosis; Arteriovenous fistula thrombosis; Arteriovenous graft site stenosis; Arteriovenous graft thrombosis; Arteritis; Arteritis

090177e196ea1800\Approved\Approved On: 30-Apr-2021 09:26 (GMT)

coronary; Arthralgia; Arthritis; Arthritis enteropathic; Ascites; Aseptic cavernous sinus thrombosis; Aspartate aminotransferase abnormal; Aspartate aminotransferase increased; Aspartate-glutamate-transporter deficiency; AST to platelet ratio index increased; AST/ALT ratio abnormal; Asthma; Asymptomatic COVID-19; Ataxia; Atheroembolism; Atonic seizures; Atrial thrombosis; Atrophic thyroiditis; Atypical benign partial epilepsy; Atypical pneumonia; Aura; Autoantibody positive; Autoimmune anaemia; Autoimmune aplastic anaemia; Autoimmune arthritis; Autoimmune blistering disease; Autoimmune cholangitis; Autoimmune colitis; Autoimmune demyelinating disease; Autoimmune dermatitis; Autoimmune disorder; Autoimmune encephalopathy; Autoimmune endocrine disorder; Autoimmune enteropathy; Autoimmune eye disorder; Autoimmune haemolytic anaemia; Autoimmune heparin-induced thrombocytopenia; Autoimmune hepatitis; Autoimmune hyperlipidaemia; Autoimmune hypothyroidism; Autoimmune inner ear disease; Autoimmune lung disease; Autoimmune lymphoproliferative syndrome; Autoimmune myocarditis; Autoimmune myositis; Autoimmune nephritis; Autoimmune neuropathy; Autoimmune neutropenia; Autoimmune pancreatitis; Autoimmune pancytopenia; Autoimmune pericarditis; Autoimmune retinopathy; Autoimmune thyroid disorder; Autoimmune thyroiditis; Autoimmune uveitis; Autoinflammation with infantile enterocolitis; Autoinflammatory disease; Automatism epileptic; Autonomic nervous system imbalance; Autonomic seizure; Axial spondyloarthritis; Axillary vein thrombosis; Axonal and demyelinating polyneuropathy; Axonal neuropathy; Bacterascites; Baltic myoclonic epilepsy; Band sensation; Basedow's disease; Basilar artery thrombosis; Basophilopenia; B-cell aplasia; Behcet's syndrome; Benign ethnic neutropenia; Benign familial neonatal convulsions; Benign familial pemphigus; Benign rolandic epilepsy; Beta-2 glycoprotein antibody positive; Bickerstaff's encephalitis; Bile output abnormal; Bile output decreased; Biliary ascites; Bilirubin conjugated abnormal; Bilirubin conjugated increased; Bilirubin urine present; Biopsy liver abnormal; Biotinidase deficiency; Birdshot chorioretinopathy; Blood alkaline phosphatase abnormal; Blood alkaline phosphatase increased; Blood bilirubin abnormal; Blood bilirubin increased; Blood bilirubin unconjugated increased; Blood cholinesterase abnormal; Blood cholinesterase decreased; Blood pressure decreased; Blood pressure diastolic decreased; Blood pressure systolic decreased; Blue toe syndrome; Brachiocephalic vein thrombosis; Brain stem embolism; Brain stem thrombosis; Bromosulphthalein test abnormal; Bronchial oedema; Bronchitis; Bronchitis mycoplasmal; Bronchitis viral; Bronchopulmonary aspergillosis allergic; Bronchospasm; Budd-Chiari syndrome; Bulbar palsy; Butterfly rash; C1q nephropathy; Caesarean section; Calcium embolism; Capillaritis; Caplan's syndrome; Cardiac amyloidosis; Cardiac arrest; Cardiac failure; Cardiac failure acute; Cardiac sarcoidosis; Cardiac ventricular thrombosis; Cardiogenic shock; Cardiolipin antibody positive; Cardiopulmonary failure; Cardio-respiratory arrest; Cardio-respiratory distress; Cardiovascular insufficiency; Carotid arterial embolus; Carotid artery thrombosis; Cataplexy; Catheter site thrombosis; Catheter site vasculitis; Cavernous sinus thrombosis; CDKL5 deficiency disorder; CEC syndrome; Cement embolism; Central nervous system lupus; Central nervous system vasculitis; Cerebellar artery thrombosis; Cerebellar embolism; Cerebral amyloid angiopathy; Cerebral arteritis; Cerebral artery embolism; Cerebral artery thrombosis; Cerebral gas embolism; Cerebral microembolism; Cerebral septic infarct; Cerebral thrombosis; Cerebral venous sinus thrombosis; Cerebral venous thrombosis; Cerebrospinal thrombotic

tamponade; Cerebrovascular accident; Change in seizure presentation; Chest discomfort; Child-Pugh-Turcotte score abnormal; Child-Pugh-Turcotte score increased; Chillblains; Choking; Choking sensation; Cholangitis sclerosing; Chronic autoimmune glomerulonephritis; Chronic cutaneous lupus erythematosus; Chronic fatigue syndrome; Chronic gastritis; Chronic inflammatory demyelinating polyradiculoneuropathy; Chronic lymphocytic inflammation with pontine perivascular enhancement responsive to steroids; Chronic recurrent multifocal osteomyelitis; Chronic respiratory failure; Chronic spontaneous urticaria; Circulatory collapse; Circumoral oedema; Circumoral swelling; Clinically isolated syndrome; Clonic convulsion; Coeliac disease; Cogan's syndrome; Cold agglutinins positive; Cold type haemolytic anaemia; Colitis; Colitis erosive; Colitis herpes; Colitis microscopic; Colitis ulcerative; Collagen disorder; Collagen-vascular disease; Complement factor abnormal; Complement factor C1 decreased; Complement factor C2 decreased; Complement factor C3 decreased; Complement factor C4 decreased; Complement factor decreased; Computerised tomogram liver abnormal; Concentric sclerosis; Congenital anomaly; Congenital bilateral perisylvian syndrome; Congenital herpes simplex infection; Congenital myasthenic syndrome; Congenital varicella infection; Congestive hepatopathy; Convulsion in childhood; Convulsions local; Convulsive threshold lowered; Coombs positive haemolytic anaemia; Coronary artery disease; Coronary artery embolism; Coronary artery thrombosis; Coronary bypass thrombosis; Coronavirus infection; Coronavirus test; Coronavirus test negative; Coronavirus test positive; Corpus callosotomy; Cough; Cough variant asthma; COVID-19; COVID-19 immunisation; COVID-19 pneumonia; COVID-19 prophylaxis; COVID-19 treatment; Cranial nerve disorder; Cranial nerve palsies multiple; Cranial nerve paralysis; CREST syndrome; Crohn's disease; Cryofibrinogenaemia; Cryoglobulinaemia; CSF oligoclonal band present; CSWS syndrome; Cutaneous amyloidosis; Cutaneous lupus erythematosus; Cutaneous sarcoidosis; Cutaneous vasculitis; Cyanosis; Cyclic neutropenia; Cystitis interstitial; Cytokine release syndrome; Cytokine storm; De novo purine synthesis inhibitors associated acute inflammatory syndrome; Death neonatal; Deep vein thrombosis; Deep vein thrombosis postoperative; Deficiency of bile secretion; Deja vu; Demyelinating polyneuropathy; Demyelination; Dermatitis; Dermatitis bullous; Dermatitis herpetiformis; Dermatomyositis; Device embolisation; Device related thrombosis; Diabetes mellitus; Diabetic ketoacidosis; Diabetic mastopathy; Dialysis amyloidosis; Dialysis membrane reaction; Diastolic hypotension; Diffuse vasculitis; Digital pitting scar; Disseminated intravascular coagulation; Disseminated intravascular coagulation in newborn; Disseminated neonatal herpes simplex; Disseminated varicella; Disseminated varicella zoster vaccine virus infection; Disseminated varicella zoster virus infection; DNA antibody positive; Double cortex syndrome; Double stranded DNA antibody positive; Dreamy state; Dressler's syndrome; Drop attacks; Drug withdrawal convulsions; Dyspnoea; Early infantile epileptic encephalopathy with burst-suppression; Eclampsia; Eczema herpeticum; Embolia cutis medicamentosa; Embolic cerebellar infarction; Embolic cerebral infarction; Embolic pneumonia; Embolic stroke; Embolism; Embolism arterial; Embolism venous; Encephalitis; Encephalitis allergic; Encephalitis autoimmune; Encephalitis brain stem; Encephalitis haemorrhagic; Encephalitis periaxialis diffusa; Encephalitis post immunisation; Encephalomyelitis; Encephalopathy; Endocrine disorder; Endocrine ophthalmopathy; Endotracheal intubation; Enteritis; Enteritis leukopenic; Enterobacter pneumonia; Enterocolitis; Enteropathic spondylitis; Eosinopenia; Eosinophilic

fasciitis;Eosinophilic granulomatosis with polyangiitis;Eosinophilic oesophagitis;Epidermolysis;Epilepsy;Epilepsy surgery;Epilepsy with myoclonic-atonic seizures;Epileptic aura;Epileptic psychosis;Erythema;Erythema induratum;Erythema multiforme;Erythema nodosum;Evans syndrome;Exanthema subitum;Expanded disability status scale score decreased;Expanded disability status scale score increased;Exposure to communicable disease;Exposure to SARS-CoV-2;Eye oedema;Eye pruritus;Eye swelling;Eyelid oedema;Face oedema;Facial paralysis;Facial paresis;Faciobrachial dystonic seizure;Fat embolism;Febrile convulsion;Febrile infection-related epilepsy syndrome;Febrile neutropenia;Felty's syndrome;Femoral artery embolism;Fibrillary glomerulonephritis;Fibromyalgia;Flushing;Foaming at mouth;Focal cortical resection;Focal dyscognitive seizures;Foetal distress syndrome;Foetal placental thrombosis;Foetor hepaticus;Foreign body embolism;Frontal lobe epilepsy;Fulminant type 1 diabetes mellitus;Galactose elimination capacity test abnormal;Galactose elimination capacity test decreased;Gamma-glutamyltransferase abnormal;Gamma-glutamyltransferase increased;Gastritis herpes;Gastrointestinal amyloidosis;Gelastic seizure;Generalised onset non-motor seizure;Generalised tonic-clonic seizure;Genital herpes;Genital herpes simplex;Genital herpes zoster;Giant cell arteritis;Glomerulonephritis;Glomerulonephritis membranoproliferative;Glomerulonephritis membranous;Glomerulonephritis rapidly progressive;Glossopharyngeal nerve paralysis;Glucose transporter type 1 deficiency syndrome;Glutamate dehydrogenase increased;Glycocholic acid increased;GM2 gangliosidosis;Goodpasture's syndrome;Graft thrombosis;Granulocytopenia;Granulocytopenia neonatal;Granulomatosis with polyangiitis;Granulomatous dermatitis;Grey matter heterotopia;Guanase increased;Guillain-Barre syndrome;Haemolytic anaemia;Haemophagocytic lymphohistiocytosis;Haemorrhage;Haemorrhagic ascites;Haemorrhagic disorder;Haemorrhagic pneumonia;Haemorrhagic varicella syndrome;Haemorrhagic vasculitis;Hantavirus pulmonary infection;Hashimoto's encephalopathy;Hashitoxicosis;Hemimegalencephaly;Henoch-Schonlein purpura;Henoch-Schonlein purpura nephritis;Hepaplastin abnormal;Hepaplastin decreased;Heparin-induced thrombocytopenia;Hepatic amyloidosis;Hepatic artery embolism;Hepatic artery flow decreased;Hepatic artery thrombosis;Hepatic enzyme abnormal;Hepatic enzyme decreased;Hepatic enzyme increased;Hepatic fibrosis marker abnormal;Hepatic fibrosis marker increased;Hepatic function abnormal;Hepatic hydrothorax;Hepatic hypertrophy;Hepatic hypoperfusion;Hepatic lymphocytic infiltration;Hepatic mass;Hepatic pain;Hepatic sequestration;Hepatic vascular resistance increased;Hepatic vascular thrombosis;Hepatic vein embolism;Hepatic vein thrombosis;Hepatic venous pressure gradient abnormal;Hepatic venous pressure gradient increased;Hepatitis;Hepatobiliary scan abnormal;Hepatomegaly;Hepatosplenomegaly;Hereditary angioedema with C1 esterase inhibitor deficiency;Herpes dermatitis;Herpes gestationis;Herpes oesophagitis;Herpes ophthalmic;Herpes pharyngitis;Herpes sepsis;Herpes simplex;Herpes simplex cervicitis;Herpes simplex colitis;Herpes simplex encephalitis;Herpes simplex gastritis;Herpes simplex hepatitis;Herpes simplex meningitis;Herpes simplex meningoencephalitis;Herpes simplex meningomyelitis;Herpes simplex necrotising retinopathy;Herpes simplex oesophagitis;Herpes simplex otitis externa;Herpes simplex pharyngitis;Herpes simplex pneumonia;Herpes simplex reactivation;Herpes simplex sepsis;Herpes simplex viraemia;Herpes simplex virus conjunctivitis neonatal;Herpes simplex visceral;Herpes virus

infection;Herpes zoster;Herpes zoster cutaneous disseminated;Herpes zoster infection neurological;Herpes zoster meningitis;Herpes zoster meningoencephalitis;Herpes zoster meningomyelitis;Herpes zoster meningoradiculitis;Herpes zoster necrotising retinopathy;Herpes zoster oticus;Herpes zoster pharyngitis;Herpes zoster reactivation;Herpetic radiculopathy;Histone antibody positive;Hoigne's syndrome;Human herpesvirus 6 encephalitis;Human herpesvirus 6 infection;Human herpesvirus 6 infection reactivation;Human herpesvirus 7 infection;Human herpesvirus 8 infection;Hyperammonaemia;Hyperbilirubinaemia;Hypercholia;Hypergammaglobulinaemia benign monoclonal;Hyperglycaemic seizure;Hypersensitivity;Hypersensitivity vasculitis;Hyperthyroidism;Hypertransaminaemia;Hyperventilation;Hypoalbuminaemia;Hypocalcaemic seizure;Hypogammaglobulinaemia;Hypoglossal nerve paralysis;Hypoglossal nerve paresis;Hypoglycaemic seizure;Hyponatraemic seizure;Hypotension;Hypotensive crisis;Hypothenar hammer syndrome;Hypothyroidism;Hypoxia;Idiopathic CD4 lymphocytopenia;Idiopathic generalised epilepsy;Idiopathic interstitial pneumonia;Idiopathic neutropenia;Idiopathic pulmonary fibrosis;IgA nephropathy;IgM nephropathy;IIIrd nerve paralysis;IIIrd nerve paresis;Iliac artery embolism;Immune thrombocytopenia;Immune-mediated adverse reaction;Immune-mediated cholangitis;Immune-mediated cholestasis;Immune-mediated cytopenia;Immune-mediated encephalitis;Immune-mediated encephalopathy;Immune-mediated endocrinopathy;Immune-mediated enterocolitis;Immune-mediated gastritis;Immune-mediated hepatic disorder;Immune-mediated hepatitis;Immune-mediated hyperthyroidism;Immune-mediated hypothyroidism;Immune-mediated myocarditis;Immune-mediated myositis;Immune-mediated nephritis;Immune-mediated neuropathy;Immune-mediated pancreatitis;Immune-mediated pneumonitis;Immune-mediated renal disorder;Immune-mediated thyroiditis;Immune-mediated uveitis;Immunoglobulin G4 related disease;Immunoglobulins abnormal;Implant site thrombosis;Inclusion body myositis;Infantile genetic agranulocytosis;Infantile spasms;Infected vasculitis;Infective thrombosis;Inflammation;Inflammatory bowel disease;Infusion site thrombosis;Infusion site vasculitis;Injection site thrombosis;Injection site urticaria;Injection site vasculitis;Instillation site thrombosis;Insulin autoimmune syndrome;Interstitial granulomatous dermatitis;Interstitial lung disease;Intracardiac mass;Intracardiac thrombus;Intracranial pressure increased;Intrapericardial thrombosis;Intrinsic factor antibody abnormal;Intrinsic factor antibody positive;IPEX syndrome;Irregular breathing;IRVAN syndrome;IVth nerve paralysis;IVth nerve paresis;JC polyomavirus test positive;JC virus CSF test positive;Jeavons syndrome;Jugular vein embolism;Jugular vein thrombosis;Juvenile idiopathic arthritis;Juvenile myoclonic epilepsy;Juvenile polymyositis;Juvenile psoriatic arthritis;Juvenile spondyloarthritis;Kaposi sarcoma inflammatory cytokine syndrome;Kawasaki's disease;Kayser-Fleischer ring;Keratoderma blenorrhagica;Ketosis-prone diabetes mellitus;Kounis syndrome;Lafora's myoclonic epilepsy;Lambli's excrescences;Laryngeal dyspnoea;Laryngeal oedema;Laryngeal rheumatoid arthritis;Laryngospasm;Laryngotracheal oedema;Latent autoimmune diabetes in adults;LE cells present;Lemierre syndrome;Lennox-Gastaut syndrome;Leucine aminopeptidase increased;Leukoencephalomyelitis;Leukoencephalopathy;Leukopenia;Leukopenia neonatal;Lewis-Sumner syndrome;Lhermitte's sign;Lichen planopilaris;Lichen planus;Lichen sclerosus;Limbic encephalitis;Linear IgA disease;Lip oedema;Lip swelling;Liver function test abnormal;Liver function test decreased;Liver function test increased;Liver induration;Liver injury;Liver iron concentration abnormal;Liver iron concentration

increased;Liver opacity;Liver palpable;Liver sarcoidosis;Liver scan abnormal;Liver tenderness;Low birth weight baby;Lower respiratory tract herpes infection;Lower respiratory tract infection;Lower respiratory tract infection viral;Lung abscess;Lupoid hepatic cirrhosis;Lupus cystitis;Lupus encephalitis;Lupus endocarditis;Lupus enteritis;Lupus hepatitis;Lupus myocarditis;Lupus myositis;Lupus nephritis;Lupus pancreatitis;Lupus pleurisy;Lupus pneumonitis;Lupus vasculitis;Lupus-like syndrome;Lymphocytic hypophysitis;Lymphocytopenia neonatal;Lymphopenia;MAGIC syndrome;Magnetic resonance imaging liver abnormal;Magnetic resonance proton density fat fraction measurement;Mahler sign;Manufacturing laboratory analytical testing issue;Manufacturing materials issue;Manufacturing production issue;Marburg's variant multiple sclerosis;Marchiafava-Bignami disease;Marine Lenhart syndrome;Mastocytic enterocolitis;Maternal exposure during pregnancy;Medical device site thrombosis;Medical device site vasculitis;MELAS syndrome;Meningitis;Meningitis aseptic;Meningitis herpes;Meningoencephalitis herpes simplex neonatal;Meningoencephalitis herpetic;Meningomyelitis herpes;MERS-CoV test;MERS-CoV test negative;MERS-CoV test positive;Mesangioproliferative glomerulonephritis;Mesenteric artery embolism;Mesenteric artery thrombosis;Mesenteric vein thrombosis;Metapneumovirus infection;Metastatic cutaneous Crohn's disease;Metastatic pulmonary embolism;Microangiopathy;Microembolism;Microscopic polyangiitis;Middle East respiratory syndrome;Migraine-triggered seizure;Miliary pneumonia;Miller Fisher syndrome;Mitochondrial aspartate aminotransferase increased;Mixed connective tissue disease;Model for end stage liver disease score abnormal;Model for end stage liver disease score increased;Molar ratio of total branched-chain amino acid to tyrosine;Molybdenum cofactor deficiency;Monocytopenia;Mononeuritis;Mononeuropathy multiplex;Morphoea;Morvan syndrome;Mouth swelling;Moyamoya disease;Multifocal motor neuropathy;Multiple organ dysfunction syndrome;Multiple sclerosis;Multiple sclerosis relapse;Multiple sclerosis relapse prophylaxis;Multiple subpial transection;Multisystem inflammatory syndrome in children;Muscular sarcoidosis;Myasthenia gravis;Myasthenia gravis crisis;Myasthenia gravis neonatal;Myasthenic syndrome;Myelitis;Myelitis transverse;Myocardial infarction;Myocarditis;Myocarditis post infection;Myoclonic epilepsy;Myoclonic epilepsy and ragged-red fibres;Myokymia;Myositis;Narcolepsy;Nasal herpes;Nasal obstruction;Necrotising herpetic retinopathy;Neonatal Crohn's disease;Neonatal epileptic seizure;Neonatal lupus erythematosus;Neonatal mucocutaneous herpes simplex;Neonatal pneumonia;Neonatal seizure;Nephritis;Nephrogenic systemic fibrosis;Neuralgic amyotrophy;Neuritis;Neuritis cranial;Neuromyelitis optica pseudo relapse;Neuromyelitis optica spectrum disorder;Neuromyotonia;Neuronal neuropathy;Neuropathy peripheral;Neuropathy, ataxia, retinitis pigmentosa syndrome;Neuropsychiatric lupus;Neurosarcoidosis;Neutropenia;Neutropenia neonatal;Neutropenic colitis;Neutropenic infection;Neutropenic sepsis;Nodular rash;Nodular vasculitis;Noninfectious myelitis;Noninfective encephalitis;Noninfective encephalomyelitis;Noninfective oophoritis;Obstetrical pulmonary embolism;Occupational exposure to communicable disease;Occupational exposure to SARS-CoV-2;Ocular hyperaemia;Ocular myasthenia;Ocular pemphigoid;Ocular sarcoidosis;Ocular vasculitis;Oculofacial paralysis;Oedema;Oedema blister;Oedema due to hepatic disease;Oedema mouth;Oesophageal achalasia;Ophthalmic artery thrombosis;Ophthalmic herpes simplex;Ophthalmic herpes zoster;Ophthalmic vein thrombosis;Optic neuritis;Optic

neuropathy; Optic perineuritis; Oral herpes; Oral lichen planus; Oropharyngeal oedema; Oropharyngeal spasm; Oropharyngeal swelling; Osmotic demyelination syndrome; Ovarian vein thrombosis; Overlap syndrome; Paediatric autoimmune neuropsychiatric disorders associated with streptococcal infection; Paget-Schroetter syndrome; Palindromic rheumatism; Palisaded neutrophilic granulomatous dermatitis; Palmoplantar keratoderma; Palpable purpura; Pancreatitis; Panencephalitis; Papillophlebitis; Paracancerous pneumonia; Paradoxical embolism; Parainfluenzae viral laryngotracheobronchitis; Paraneoplastic dermatomyositis; Paraneoplastic pemphigus; Paraneoplastic thrombosis; Paresis cranial nerve; Parietal cell antibody positive; Paroxysmal nocturnal haemoglobinuria; Partial seizures; Partial seizures with secondary generalisation; Patient isolation; Pelvic venous thrombosis; Pemphigoid; Pemphigus; Penile vein thrombosis; Pericarditis; Pericarditis lupus; Perihepatic discomfort; Periorbital oedema; Periorbital swelling; Peripheral artery thrombosis; Peripheral embolism; Peripheral ischaemia; Peripheral vein thrombus extension; Periportal oedema; Peritoneal fluid protein abnormal; Peritoneal fluid protein decreased; Peritoneal fluid protein increased; Peritonitis lupus; Pernicious anaemia; Petit mal epilepsy; Pharyngeal oedema; Pharyngeal swelling; Pityriasis lichenoides et varioliformis acuta; Placenta praevia; Pleuroparenchymal fibroelastosis; Pneumobilia; Pneumonia; Pneumonia adenoviral; Pneumonia cytomegaloviral; Pneumonia herpes viral; Pneumonia influenzal; Pneumonia measles; Pneumonia mycoplasmal; Pneumonia necrotising; Pneumonia parainfluenzae viral; Pneumonia respiratory syncytial viral; Pneumonia viral; POEMS syndrome; Polyarteritis nodosa; Polyarthritis; Polychondritis; Polyglandular autoimmune syndrome type I; Polyglandular autoimmune syndrome type II; Polyglandular autoimmune syndrome type III; Polyglandular disorder; Polymicrogyria; Polymyalgia rheumatica; Polymyositis; Polyneuropathy; Polyneuropathy idiopathic progressive; Portal pyaemia; Portal vein embolism; Portal vein flow decreased; Portal vein pressure increased; Portal vein thrombosis; Portosplenomesenteric venous thrombosis; Post procedural hypotension; Post procedural pneumonia; Post procedural pulmonary embolism; Post stroke epilepsy; Post stroke seizure; Post thrombotic retinopathy; Post thrombotic syndrome; Post viral fatigue syndrome; Postictal headache; Postictal paralysis; Postictal psychosis; Postictal state; Postoperative respiratory distress; Postoperative respiratory failure; Postoperative thrombosis; Postpartum thrombosis; Postpartum venous thrombosis; Postpericardiotomy syndrome; Post-traumatic epilepsy; Postural orthostatic tachycardia syndrome; Precerebral artery thrombosis; Pre-eclampsia; Preictal state; Premature labour; Premature menopause; Primary amyloidosis; Primary biliary cholangitis; Primary progressive multiple sclerosis; Procedural shock; Proctitis herpes; Proctitis ulcerative; Product availability issue; Product distribution issue; Product supply issue; Progressive facial hemiatrophy; Progressive multifocal leukoencephalopathy; Progressive multiple sclerosis; Progressive relapsing multiple sclerosis; Prosthetic cardiac valve thrombosis; Pruritus; Pruritus allergic; Pseudovasculitis; Psoriasis; Psoriatic arthropathy; Pulmonary amyloidosis; Pulmonary artery thrombosis; Pulmonary embolism; Pulmonary fibrosis; Pulmonary haemorrhage; Pulmonary microemboli; Pulmonary oil microembolism; Pulmonary renal syndrome; Pulmonary sarcoidosis; Pulmonary sepsis; Pulmonary thrombosis; Pulmonary tumour thrombotic microangiopathy; Pulmonary vasculitis; Pulmonary veno-occlusive disease; Pulmonary venous thrombosis; Pyoderma gangrenosum; Pyostomatitis vegetans; Pyrexia; Quarantine; Radiation leukopenia; Radiculitis

brachial;Radiologically isolated syndrome;Rash;Rash erythematous;Rash pruritic;Rasmussen encephalitis;Raynaud's phenomenon;Reactive capillary endothelial proliferation;Relapsing multiple sclerosis;Relapsing-remitting multiple sclerosis;Renal amyloidosis;Renal arteritis;Renal artery thrombosis;Renal embolism;Renal failure;Renal vascular thrombosis;Renal vasculitis;Renal vein embolism;Renal vein thrombosis;Respiratory arrest;Respiratory disorder;Respiratory distress;Respiratory failure;Respiratory paralysis;Respiratory syncytial virus bronchiolitis;Respiratory syncytial virus bronchitis;Retinal artery embolism;Retinal artery occlusion;Retinal artery thrombosis;Retinal vascular thrombosis;Retinal vasculitis;Retinal vein occlusion;Retinal vein thrombosis;Retinol binding protein decreased;Retinopathy;Retrograde portal vein flow;Retroperitoneal fibrosis;Reversible airways obstruction;Reynold's syndrome;Rheumatic brain disease;Rheumatic disorder;Rheumatoid arthritis;Rheumatoid factor increased;Rheumatoid factor positive;Rheumatoid factor quantitative increased;Rheumatoid lung;Rheumatoid neutrophilic dermatosis;Rheumatoid nodule;Rheumatoid nodule removal;Rheumatoid scleritis;Rheumatoid vasculitis;Saccadic eye movement;SAPHO syndrome;Sarcoidosis;SARS-CoV-1 test;SARS-CoV-1 test negative;SARS-CoV-1 test positive;SARS-CoV-2 antibody test;SARS-CoV-2 antibody test negative;SARS-CoV-2 antibody test positive;SARS-CoV-2 carrier;SARS-CoV-2 sepsis;SARS-CoV-2 test;SARS-CoV-2 test false negative;SARS-CoV-2 test false positive;SARS-CoV-2 test negative;SARS-CoV-2 test positive;SARS-CoV-2 viraemia;Satoyoshi syndrome;Schizencephaly;Scleritis;Sclerodactylia;Scleroderma;Scleroderma associated digital ulcer;Scleroderma renal crisis;Scleroderma-like reaction;Secondary amyloidosis;Secondary cerebellar degeneration;Secondary progressive multiple sclerosis;Segmented hyalinising vasculitis;Seizure;Seizure anoxic;Seizure cluster;Seizure like phenomena;Seizure prophylaxis;Sensation of foreign body;Septic embolus;Septic pulmonary embolism;Severe acute respiratory syndrome;Severe myoclonic epilepsy of infancy;Shock;Shock symptom;Shrinking lung syndrome;Shunt thrombosis;Silent thyroiditis;Simple partial seizures;Sjogren's syndrome;Skin swelling;SLE arthritis;Smooth muscle antibody positive;Sneezing;Spinal artery embolism;Spinal artery thrombosis;Splenic artery thrombosis;Splenic embolism;Splenic thrombosis;Splenic vein thrombosis;Spondylitis;Spondyloarthropathy;Spontaneous heparin-induced thrombocytopenia syndrome;Status epilepticus;Stevens-Johnson syndrome;Stiff leg syndrome;Stiff person syndrome;Stillbirth;Still's disease;Stoma site thrombosis;Stoma site vasculitis;Stress cardiomyopathy;Stridor;Subacute cutaneous lupus erythematosus;Subacute endocarditis;Subacute inflammatory demyelinating polyneuropathy;Subclavian artery embolism;Subclavian artery thrombosis;Subclavian vein thrombosis;Sudden unexplained death in epilepsy;Superior sagittal sinus thrombosis;Susac's syndrome;Suspected COVID-19;Swelling;Swelling face;Swelling of eyelid;Swollen tongue;Sympathetic ophthalmia;Systemic lupus erythematosus;Systemic lupus erythematosus disease activity index abnormal;Systemic lupus erythematosus disease activity index decreased;Systemic lupus erythematosus disease activity index increased;Systemic lupus erythematosus rash;Systemic scleroderma;Systemic sclerosis pulmonary;Tachycardia;Tachypnoea;Takayasu's arteritis;Temporal lobe epilepsy;Terminal ileitis;Testicular autoimmunity;Throat tightness;Thromboangiitis obliterans;Thrombocytopenia;Thrombocytopenic purpura;Thrombophlebitis;Thrombophlebitis migrans;Thrombophlebitis

neonatal;Thrombophlebitis septic;Thrombophlebitis superficial;Thromboplastin antibody positive;Thrombosis;Thrombosis corpora cavernosa;Thrombosis in device;Thrombosis mesenteric vessel;Thrombotic cerebral infarction;Thrombotic microangiopathy;Thrombotic stroke;Thrombotic thrombocytopenic purpura;Thyroid disorder;Thyroid stimulating immunoglobulin increased;Thyroiditis;Tongue amyloidosis;Tongue biting;Tongue oedema;Tonic clonic movements;Tonic convulsion;Tonic posturing;Topectomy;Total bile acids increased;Toxic epidermal necrolysis;Toxic leukoencephalopathy;Toxic oil syndrome;Tracheal obstruction;Tracheal oedema;Tracheobronchitis;Tracheobronchitis mycoplasmal;Tracheobronchitis viral;Transaminases abnormal;Transaminases increased;Transfusion-related alloimmune neutropenia;Transient epileptic amnesia;Transverse sinus thrombosis;Trigeminal nerve paresis;Trigeminal neuralgia;Trigeminal palsy;Truncus coeliacus thrombosis;Tuberous sclerosis complex;Tubulointerstitial nephritis and uveitis syndrome;Tumefactive multiple sclerosis;Tumour embolism;Tumour thrombosis;Type 1 diabetes mellitus;Type I hypersensitivity;Type III immune complex mediated reaction;Uhthoff's phenomenon;Ulcerative keratitis;Ultrasound liver abnormal;Umbilical cord thrombosis;Uncinate fits;Undifferentiated connective tissue disease;Upper airway obstruction;Urine bilirubin increased;Urobilinogen urine decreased;Urobilinogen urine increased;Urticaria;Urticaria papular;Urticarial vasculitis;Uterine rupture;Uveitis;Vaccination site thrombosis;Vaccination site vasculitis;Vagus nerve paralysis;Varicella;Varicella keratitis;Varicella post vaccine;Varicella zoster gastritis;Varicella zoster oesophagitis;Varicella zoster pneumonia;Varicella zoster sepsis;Varicella zoster virus infection;Vasa praevia;Vascular graft thrombosis;Vascular pseudoaneurysm thrombosis;Vascular purpura;Vascular stent thrombosis;Vasculitic rash;Vasculitic ulcer;Vasculitis;Vasculitis gastrointestinal;Vasculitis necrotising;Vena cava embolism;Vena cava thrombosis;Venous intravasation;Venous recanalisation;Venous thrombosis;Venous thrombosis in pregnancy;Venous thrombosis limb;Venous thrombosis neonatal;Vertebral artery thrombosis;Vessel puncture site thrombosis;Visceral venous thrombosis;VIth nerve paralysis;VIth nerve paresis;Vitiligo;Vocal cord paralysis;Vocal cord paresis;Vogt-Koyanagi-Harada disease;Warm type haemolytic anaemia;Wheezing;White nipple sign;XIth nerve paralysis;X-ray hepatobiliary abnormal;Young's syndrome;Zika virus associated Guillain Barre syndrome.